

Managing Children who are Sick, Infectious, or with Allergies

November 2023

Policy statement

We aim to provide care for healthy children through preventing cross infection of viruses and bacterial infections and promote health through identifying allergies and preventing contact with the allergenic trigger.

Procedures for children who are sick or infectious

- If a child appears unwell during the day, for example has a raised temperature, sickness, diarrhoea* and/or pains, particularly in the head or stomach then the setting manager calls the parents and asks them to collect the child or send a known carer to collect on their behalf.
- If a child has a raised temperature, they are kept cool by removing top clothing, sponging their heads with cool water and kept away from draughts.
- A child's temperature is taken and checked regularly, using Fever Scans or other means i.e. ear thermometer.
- In an emergency an ambulance is called and the parents are informed.
- If a child has a worryingly high temperature, and it is not dropping, then parental consent will be sought so that Calpol (or other branded paracetamol medication) can be administered, to reduce the risk of Fibril convulsions, whilst the child is waiting to be collected.
- Parents are advised to seek medical advice before returning them to the setting; the setting can refuse admittance to children who have a raised temperature, sickness and diarrhoea or a contagious infection or disease.

- Where children have been prescribed antibiotics for an infectious illness or complaint, parents are asked to keep them at home for 48 hours.
- After diarrhoea or vomiting, parents are asked to keep children home for 48 hours following the last episode.
- Some activities such as sand and water play and self-serve snack will be suspended for the duration of any outbreak.
- The setting has information about excludable diseases and exclusion times.
- The setting manager notifies their line manager if there is an outbreak of an infection (affects more than 3-4 children) and keeps a record of the numbers and duration of each event.
- The setting manager has a list of notifiable diseases and contacts Public Health England (PHE) and Ofsted in the event of an outbreak.
- If staff suspect that a child who falls ill whilst in their care is suffering from a serious disease that may have been contracted abroad such as Ebola, immediate medical assessment is required. The setting manager or deputy calls NHS111 and informs parents.

HIV/AIDS procedure

HIV virus, like other viruses such as Hepatitis, (A, B and C), are spread through body fluids. Hygiene precautions for dealing with body fluids are the same for all children and adults.

- Single use vinyl gloves and aprons are worn when changing children's nappies, pants and clothing that are soiled with blood, urine, faeces or vomit.
- Protective rubber gloves are used for cleaning/slucing clothing after changing.
- Soiled clothing is rinsed and bagged for parents to collect.
- Spills of blood, urine, faeces or vomit are cleared using mild disinfectant solution and mops; cloths used are disposed of with clinical waste.
- Tables and other furniture or toys affected by blood, urine, faeces or vomit are cleaned using a disinfectant.
- Baby mouthing toys are kept clean and plastic toys cleaned in sterilising solution regularly.

Nits and head lice

- Nits and head lice are not an excludable condition; although in exceptional cases parents may be asked to keep the child away from the setting until the infestation has cleared.
- On identifying cases of head lice, all parents are informed and asked to treat their child and all the family, using current recommended treatments methods if they are found.

*Diarrhoea is defined as 3 or more liquid or semi-liquid stools in a 24-hour period.

(www.gov.uk/government/publications/health-protection-in-schools-and-other-childcare-facilities/chapter-9-managing-specific-infectious-diseases#diarrhoea-and-vomiting-gastroenteritis)

Managing a suspected case of Coronavirus

The main symptoms of coronavirus are:

- a high temperature
- a new continuous cough – this means coughing a lot, for more than an hour, or three or more coughing episodes in 24 hours
- a loss of change to smell or taste – this means they cannot smell or taste anything, or things smell or taste different to normal

Please refer to the latest government guidance on [next-steps-for-living-with-COVID](#). If it is suspected that a child has COVID, staff do not attempt to diagnose or make assumptions about symptoms presented. They should immediately respond and take action as detailed in this procedure. This includes asking parents/carers to seek further advice from a medical practitioner who may/or may not advise that the symptoms meet the criteria for testing. In which case if the child appears well and displays no further suspect symptoms, they can return to the setting within the timescale advised by the medical practitioner.

The focus on coronavirus must not detract from staff being alert to the signs and symptoms linked to other serious illness as detailed below:

What to do if a child seems very unwell

Children and babies will still get illnesses that can make them very unwell quickly. It is important to get seek medical help and to contact the child's parents immediately.

Call 999 if a child:

- has a stiff neck
- has a rash that does not fade when you press a glass against it
- is bothered by light
- has a seizure or fit for the first time
- has unusually cold hands
- has pale, blotchy, blue or grey skin
- has a weak, high-pitched cry that is not like their usual cry
- is extremely agitated (does not stop crying) or is confused
- finds it hard to breathe
- has a soft spot on their head that curves outwards
- is not responding like they normally do

Being prepared

- All staff are aware of this procedure and their responsibility if a child becomes unwell with coronavirus symptoms at the setting.
- Staff are instructed in how to remove and dispose of PPE equipment safely – this includes aprons and gloves worn during routine care procedures. We display the [NHS guide to putting on and removing PPE](#).

If a child becomes unwell

- If a child is displaying any of the symptoms of coronavirus. The manager/deputy calls their parents to collect them immediately. Current guidance states that: *'If a child or young person has a positive COVID-19 test result they should try to stay at home and where possible avoid contact with other people for 3 days after the day they took the test. The risk of passing the infection on to others is much lower after 3 days, if they feel well and do not have a high temperature. Children and young people who usually attend an education or childcare setting and who live with someone who has a positive COVID-19 test result should continue to attend as normal'*.

- We will maintain contact with the parent(s) of the child who was sent home, and ensure they know that their child is entitled to a test and encourage them to get their child tested. To access testing parents should use the [111 online coronavirus service](#).
- We will ask the parent(s) to let us know the outcome as soon as possible.
- If the test result is positive we will inform all other parents that a child has tested positive and remind them to be aware of the symptoms to look out for.
- We will inform our local authority if a child, or staff member in the setting tests positive for coronavirus.

Life-saving medication and invasive treatments

Life-saving medication and invasive treatments may include adrenaline injections (Epipens) for anaphylactic shock reactions (caused by allergies to nuts, eggs etc) or invasive treatment such as rectal administration of Diazepam (for epilepsy).

- The key person responsible for the intimate care of children who require life-saving medication or invasive treatment will undertake their duties in a professional manner having due regard to the procedures listed above.
- The child's welfare is paramount, and their experience of intimate and personal care should be positive. Every child is treated as an individual and care is given gently and sensitively; no child should be attended to in a way that causes distress or pain.
- The key person works in close partnership with parents/carers and other professionals to share information and provide continuity of care.
- Children with complex and/or long-term health conditions have a health care plan (04.2a) in place which takes into account the principles and best practice guidance given here.
- Key persons have appropriate training for administration of treatment and are aware of infection control best practice, for example, using personal protective equipment (PPE).
- Key persons speak directly to the child, explaining what they are doing as appropriate to the child's age and level of comprehension.
- Children's right to privacy and modesty is respected. Another educator is usually present during the process.

Record keeping

For a child who requires invasive treatment the following must be in place from the outset:

- a letter from the child's GP/consultant stating the child's condition and what medication if any is to be administered
- written consent from parents allowing members of staff to administer medication
- proof of training in the administration of such medication by the child's GP, a district nurse, children's nurse specialist or a community paediatric nurse
- a healthcare plan (04.2a)

Copies of all letters relating to these children must be sent to the insurance provider for appraisal. Confirmation will then be issued in writing confirming that the insurance has been extended. A record is made in the medication record book of the intimate/invasive treatment each time it is given.

Physiotherapy

- Children who require physiotherapy whilst attending the setting should have this carried out by a trained physiotherapist.
- If it is agreed in the health care plan that the key person should undertake part of the physiotherapy regime then the required technique must be demonstrated by the physiotherapist personally; written guidance must also be given and reviewed regularly. The physiotherapist should observe the educator applying the technique in the first instance.

Safeguarding/child protection

- Educators recognise that children with SEND are particularly vulnerable to all types of abuse, therefore the safeguarding procedures are followed rigorously.
- If an educator has any concerns about physical changes noted during a procedure, for example unexplained marks or bruising then the concerns are discussed with the designated person for safeguarding and the relevant procedure is followed.

Treatments such as inhalers or Epi-pens must be immediately accessible in an emergency.

Allergies and food intolerance

When a child starts at the setting, parents are asked if their child has any known allergies or food intolerance. This information is recorded on the registration form.

- If a child has an allergy or food intolerance, a **risk assessment form** is completed with the following information:
 - the risk identified – the allergen (i.e. the substance, material or living creature the child is allergic to such as nuts, eggs, bee stings, cats etc.)
 - the level of risk, taking into consideration the likelihood of the child coming into contact with the allergen
 - control measures, such as prevention from contact with the allergen
 - review measures
- A **Health care plan** form must be completed with:
 - the nature of the reaction e.g. anaphylactic shock reaction, including rash, reddening of skin, swelling, breathing problems etc.
 - managing allergic reactions, medication used and method (e.g. Epipen)
- The child's name **photograph and protocol is displayed for all staff to see in the kitchen.**
- A copy of the risk assessment and health care plan is kept in the child's personal file and is shared with all staff.
- Parents show staff how to administer medication in the event of an allergic reaction.
- Generally, no nuts or nut products are used within the setting.
- Parents are made aware, so that no nut or nut products are accidentally brought in.
- Any foods containing food allergens are identified on children's menus.

Oral Medication

- Oral medication must be prescribed or have manufacturer's instructions written on them.
- Staff must be provided with clear written instructions for administering such medication.
- All risk assessment procedures are adhered to for the correct storage and administration

of the medication.

- The setting must have the parents' prior written consent. Consent is kept on file.

Please also refer to the Administration of medicine policy.

Infection control

Good practice infection control is paramount in early years settings. Young children's immune systems are still developing, and they are therefore more susceptible to illness.

Prevention

- Minimise contact with individuals who are unwell by ensuring that those who have symptoms of an infectious illness do not attend settings and stay at home for the recommended exclusion time (see below UKHSA link).
- Always clean hands thoroughly, and more often than usual where there is an infection outbreak.
- Ensure good respiratory hygiene amongst children and staff by promoting 'catch it, bin it, kill it' approach.
- Where necessary, for instance, where there is an infection outbreak, wear appropriate PPE.

Response to an infection outbreak

- Manage confirmed cases of a contagious illness by following the guidance from the [UK Health Security Agency \(UKHSA\)](#)

Informing others

Early years providers have a duty to inform Ofsted of any serious accidents, illnesses or injuries as follows:

- anything that requires resuscitation
- admittance to hospital for more than 24 hours

- a broken bone or fracture
- dislocation of any major joint, such as the shoulder, knee, hip or elbow
- any loss of consciousness
- severe breathing difficulties, including asphyxia
- anything leading to hypothermia or heat-induced illness

In some circumstances this may include a confirmed case of a Notifiable Disease in their setting, if it meets the criteria defined by Ofsted above. Please note that it is not the responsibility of the setting to diagnose a notifiable disease. This can only be done by a clinician (GP or Doctor). If a child is displaying symptoms that indicate they may be suffering from a notifiable disease, parents must be advised to seek a medical diagnosis, which will then be 'notified' to the relevant body. Once a diagnosis is confirmed, the setting may be contacted by the UKHSA, or may wish to contact them for further advice.

Further guidance

[Good Practice in Early Years Infection Control](#) (Alliance 2009)

[Medication Administration Record](#) (Early Years Alliance 2019)

Guidance on infection control in schools and other childcare settings (Public Health Agency)

https://www.publichealth.hscni.net/sites/default/files/Guidance_on_infection_control_in%20schools_poster.pdf

[Good Practice in Early Years Infection Control](#) (Alliance 2009)

Version	Changes made	Author	Date	Review Date
1.0	Baseline version	H Clarke	November 2022	September 2023
1.1	No Changes	H Clarke	September 2023	September 2024